

# Pediatric Exanthems

By

MAJ Harry D. McKinnon Jr.

Department of Family Practice

Ft. Belvoir, VA

# Objectives

- Develop a logical approach to the evaluation of exanthems in a child
- Identify common causes of pediatric exanthems
- Understand treatment of common pediatric exanthems

# History



- Detailed history
  - Recent travel
  - Woodland exposure
  - Drug ingestion
  - Ill contacts
  - Medical history

# History

- Rash details
  - Site of onset
  - Rate, direction of spread
  - Pruritis
  - Temporal relationship of rash and fever
  - Oral or topical therapies




# Physical Examination

- Identify primary lesion and presence of secondary lesions
- Thorough examination essential to accurate diagnosis



# Laboratory Data



- Availability dependent
- As clinically appropriate
- Serologic tests not often helpful in acute setting
- Aspirates, scrapings and pustular fluid may be obtained

# Rubella (German measles)

- Etiology:  
Enveloped RNA  
Togavirus
- Transmission via  
direct contact,  
less commonly air  
droplets
- Incubation period:  
14 days



# Rubella (German measles)


- Transplacental transmission during viremia
- Pre-vaccine 90% acquired prior to age 15
- Late winter – early spring



# Rubella (German measles)



- Vaccine
  - Developed 1969
  - Live attenuated (CDC)
- Incidence
  - < 15 yrs: 0.06 per 100,000
  - 15-44 yrs: 0.24 per 100,000



# Rubella (German measles)

- Presentation
  - Prodrome uncommon in children
  - Pink macules/papules begin on forehead spread inferiorly and to extremities within 1 day
  - Fading in reverse order by 3<sup>rd</sup> day
  - Forchheimer's spots – petechiae on soft palate

# Rubella (German measles)

- Treatment:
  - Supportive



# Rubeola (measles)





- Paramyxovirus family
- Pre-vaccine incidence similar to Rubella
- Transmission via direct contact
- Incubation period: 10-12 days
- Late winter – early spring

# Rubeola (measles)

- Vaccine
  - Develop 1963
  - Live attenuated (CDC)
  - Current vaccine 95% effective, lifelong immunity
- Incidence
  - < 0.5 cases per 1,000,000 in 1999









# **Measles Resurgence - United States, 1989-1991**

- **Cases** **55,622**
- **Age group affected** **Children  $\leq$  5 yrs**
- **Hospitalizations** **>11,000**
- **Deaths** **123**
- **Direct medical costs** **>\$150 million**

# Rubeola (measles)

- Presentation
  - Prodrome of high fever, malaise, anorexia followed by URI symptoms; triad of cough, coryza and conjunctivitis
  - Appearance of macular-papular (confluent) rash on or about 4<sup>th</sup> febrile day
  - Rash first of face/neck spreads centrifugally and inferiorly; fades in 4-6 days

# Rubeola (measles)



Koplick spots  
pathognomonic

# Rubeola (measles)

- Treatment:
  - Primarily supportive
  - IgG, Ribavarin and Vitamin A may have some utility

# Varicella (Chicken pox)

- Varicella-zoster virus: member of herpes virus family, double stranded DNA viruses
- Transmission thru direct contact with respiratory secretions or lesion fluid or airborne spread
- Transplacental transmission
- Incubation period: 10-21 days



# Varicella (Chicken pox)



- Epidemiology:
  - 90% of cases in children < 10yrs
  - 5% in individuals > 15 yrs
- Treatment:
  - Supportive
  - Antivirals may have role in more significant disease

## **Varicella Clinical Features**

- **Incubation period 14-16 days (range 10-21 days)**
- **Mild prodrome for 1-2 days**
- **Generally appear first on head; most concentrated on trunk**
- **Successive crops (2-4 days) of pruritic vesicles**

# **Congenital Varicella Syndrome**

- **Results from maternal infection during pregnancy**
- **Period of risk may extend through first 20 weeks of pregnancy**
- **Atrophy of extremity with skin scarring, low birth weight, eye and neurologic abnormalities**
- **Risk appears to be small (<2%)**



# **Varicella Vaccine**

- **Composition**                      **Live virus (Oka-Merck strain)**
- **Efficacy**                              **95% (Range, 65%-100%)**
- **Duration of Immunity**              **>7 years**
- **Schedule**                              **1 Dose (<13 years of age)**

**May be administered simultaneously with measles-mumps-rubella (MMR) vaccine**

# **Breakthrough Infection**

- **Retrospective cohort study of 115,000 children vaccinated in 2 HMOs during January 1995 through December 1999**
- **Risk of breakthrough varicella 2.5 times higher if varicella vaccine administered <30 days following MMR**
- **No increased risk if varicella vaccine given simultaneously or >30 days after MMR**

**MMWR 2001;50(47):1058-61**

# **Varicella Vaccine Recommendations Adolescents and Adults**

- **Persons  $\geq 13$  years of age without history of varicella**
- **Two doses separated by 4 - 8 weeks**
- **Up to 90% of adults immune**
- **Serologic testing may be cost effective**

## **Varicella Vaccine Postexposure Prophylaxis**

- **Varicella vaccine is recommended for use in susceptible person after exposure to varicella**
  - **70%-100% effective if given within 72 hours of exposure**
  - **not effective if >5 days but will produce immunity if not infected**

# Roseola



- Aliases: exanthem subitum, roseola subitum, roseola infantalis, and sixth disease
- Etiology: Human herpes virus 6 or 7, double stranded DNA virus

# Roseola

- Common worldwide
- Self-limited benign disease
- HHV-6B primary causal agent, HHV-7 produces similar syndrome in 24-36 month olds
- Epidemiology: 6 mos to 3 yrs

# Roseola

- Presentation:
  - Sudden onset of fever lasting 1-8 days, average of 4 days (as high as 40.0 C)
  - Mild irritability and lethargy despite fevers
  - Exam may reveal cervical adenopathy (posterior cerv & occipital), tonsillar, pharyngeal and TM erythema
  - 1/3 with diarrhea & vomiting

# Roseola

- Presentation:
  - Rash appears 2-3 days following fever
  - Diffuse maculopapular eruption usually sparing face, no coalescing
  - Rash resolves 1-3 days



# Erythema infectiosum (fifth disease)

- Etiology:
  - Human Parvovirus B19
  - Smallest human DNA virus (single strand of DNA)
- Transmission
  - Respiratory secretions
  - Blood product exposure
  - Transplacental
  - Most infectious prior to exanthem

# Erythema infectiosum (fifth disease)



- Incubation period: 4-14 days
- Epidemics amongst school age children
- Primarily children between 3 to 15 years of age

# Erythema infectiosum (fifth disease)

- Presentation:

- Mild prodrome (headache, coryza, malaise) for 2-3 days prior to rash. Arthralgias, arthritis in about 10%.
- Fiery red macular rash “slapped cheeks” giving way to generalized lacy-reticular rash.
- Rash typically resolves in 5-10 days although may wax & wane for weeks or months.



# Erythema infectiosum (fifth disease)

- Treatment is supportive
- Vaccine is being developed but not available yet

# Hand, Foot and Mouth Disease



- Etiology:
  - Coxsackie A & B
  - Self-limited non-polio enterovirus
  - Highly contagious, aerosol spread
  - Bi-modal: spring and summer
  - Children < 5 yrs

# Hand, Foot and Mouth Disease

- Incubation: 4 days
- Malaise, fever, lymphadenopathy
- Oral vesicles of palate, tongue, buccal mucosa (spare gingiva) rapidly ulcerate
- Subsequent mac-pap lesions on hands & feet progress to vesicles, ulcerate then crust
- Supportive care, self-limited within 2 weeks






# Honorable mention



- Pityriasis rosea
  - Etiology unknown
  - Herald patch
  - About 5% with mild prodrome
  - Ages 10-35
  - +/- pruritis
  - Generally resolves in 2-6 weeks; may persist for months

# Recommended Childhood and Adolescent Immunization Schedule — United States, January - June 2004

Range of Recommended Ages					Catch-up Immunization				Preadolescent Assessment					
Vaccine▼	Age►	Birth	1 mo	2 mo	4 mo	6 mo	12 mo	15 mo	18 mo	24 mo	4-6 y	11-12 y	13-18 y	
Hepatitis B <sup>1</sup>		HepB #1	only if mother HBsAg (-)							HepB series				
			HepB #2			HepB #3								
Diphtheria, Tetanus, Pertussis <sup>2</sup>				DTaP	DTaP	DTaP		DTaP			DTaP	Td	Td	
<i>Haemophilus influenzae</i> Type b <sup>3</sup>				Hib	Hib	Hib <sup>3</sup>	Hib							
Inactivated Poliovirus				IPV	IPV	IPV					IPV			
Measles, Mumps, Rubella <sup>4</sup>							MMR #1				MMR #2	MMR #2		
Varicella <sup>5</sup>							Varicella			Varicella				
Pneumococcal <sup>6</sup>				PCV	PCV	PCV	PCV			PCV	PPV			
Vaccines below this line are for selected populations														
Hepatitis A <sup>7</sup>										Hepatitis A series				
Influenza <sup>8</sup>						Influenza (yearly)								

This schedule indicates the recommended ages for routine administration of currently licensed childhood vaccines, as of December 1, 2003, for children through age 18 years. Any dose not given at the recommended age should be given at any subsequent visit when indicated and feasible.  Indicates age groups that warrant special effort to administer those vaccines not previously given. Additional vaccines may be licensed and recommended during the year. Licensed combination vaccines may be used whenever any components of the combination are indicated and the vaccine's other components are not contraindicated. Providers should consult the manufacturers' package inserts for detailed recommendations. Clinically significant adverse events that follow immunization should be reported to the Vaccine Adverse Event Reporting System (VAERS). Guidance about how to obtain and complete a VAERS form can be found on the Internet: <http://www.vaers.org/> or by calling 1-800-822-7967.





